

\$%^STN;HighlightOn= \*\*\*;HighlightOff=\*\*\* ;  
Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID:sssptal65lhjg  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files  
NEWS 3 Feb 06 Engineering Information Encompass files have new names  
NEWS 4 Feb 16 TOXLINE no longer being updated  
NEWS 5 Apr 23 Search Derwent WPINDEX by chemical structure  
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA  
NEWS 7 May 07 DGENE Reload

NEWS EXPRESS May 23 CURRENT WINDOWS VERSION IS V6.0a,  
CURRENT MACINTOSH VERSION IS V5.0C (ENG) AND V5.0JB (JP),  
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2001

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 07:59:46 ON 18 JUN 2001

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.15	0.15

FILE 'REGISTRY' ENTERED AT 07:59:50 ON 18 JUN 2001  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2001 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 15 JUN 2001 HIGHEST RN 341925-25-5  
DICTIONARY FILE UPDATES: 15 JUN 2001 HIGHEST RN 341925-25-5

TSCA INFORMATION NOW CURRENT THROUGH January 11, 2001

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Structure search limits have been increased. See HELP SLIMIT  
for details.

=> e 2-pyridinol, 1-ox/cn

E1	1	2-PYRIDINOL, 1-HEXADECYL-1,2-DIHYDRO-/CN
E2	1	2-PYRIDINOL, 1-HEXADECYL-1,2-DIHYDRO-, COMPD. WITH DIBENZOYL PEROXIDE (1:1)/CN
E3	0 -->	2-PYRIDINOL, 1-OX/CN
E4	1	2-PYRIDINOL, 1-OXIDE/CN
E5	1	2-PYRIDINOL, 1-OXIDE, COMPD. WITH DODECYLAMINE/CN
E6	1	2-PYRIDINOL, 1-OXIDE, COMPD. WITH DODECYLAMINE (1:1)/CN
E7	1	2-PYRIDINOL, 1-OXIDE, ION(1-)/CN
E8	1	2-PYRIDINOL, 1-OXIDE, ION(1-), HEXADECYLTRIMETHYLAMMONIUM/CN
E9	1	2-PYRIDINOL, 1-OXIDE, LITHIUM SALT/CN
E10	1	2-PYRIDINOL, 1-OXIDE, POTASSIUM SALT/CN
E11	1	2-PYRIDINOL, 1-OXIDE, SODIUM SALT/CN
E12	1	2-PYRIDINOL, 1-OXIDE, ZINC SALT/CN

=> e

E13	1	2-PYRIDINOL, 2,4-BIS(3,5-DIMETHYLPYRROL-2-YL)-1,2-DIHYDRO-6- METHYL-/CN
E14	1	2-PYRIDINOL, 2,4-BIS(4-ETHYL-3,5-DIMETHYLPYRROL-2-YL)-1,2-DI- HYDRO-6-METHYL-/CN
E15	1	2-PYRIDINOL, 2-NITROBENZOATE (ESTER)/CN
E16	1	2-PYRIDINOL, 3,3,5,6-TETRACHLORO-2,3,4,5-TETRAHYDRO-/CN
E17	1	2-PYRIDINOL, 3,3,5,6-TETRACHLORO-2,3,4,5-TETRAHYDRO-5-METHYL -/CN
E18	1	2-PYRIDINOL, 3,3,5,6-TETRACHLORO-2,3,4,5-TETRAHYDRO-5-METHYL -, (5S)-/CN
E19	1	2-PYRIDINOL, 3,3,5,6-TETRACHLORO-2,3,4,5-TETRAHYDRO-5-METHYL -2-(TRICHLOROMETHYL)-/CN
E20	1	2-PYRIDINOL, 3,3,5,6-TETRACHLORO-2,3,4,5-TETRAHYDRO-5-METHYL -2-PHENYL-/CN
E21	1	2-PYRIDINOL, 3,4,5,6-TETRACHLORO-/CN
E22	1	2-PYRIDINOL, 3,4,5,6-TETRACHLORO-, 1-OXIDE/CN
E23	1	2-PYRIDINOL, 3,4,5,6-TETRAFLUORO-/CN
E24	1	2-PYRIDINOL, 3,4,5,6-TETRAHYDRO-, ACETATE/CN

=> s e4-e12

	1	"2-PYRIDINOL, 1-OXIDE"/CN
	1	"2-PYRIDINOL, 1-OXIDE, COMPD. WITH DODECYLAMINE"/CN
	1	"2-PYRIDINOL, 1-OXIDE, COMPD. WITH DODECYLAMINE (1:1)"/CN
	1	"2-PYRIDINOL, 1-OXIDE, ION(1-)/CN
	1	"2-PYRIDINOL, 1-OXIDE, ION(1-), HEXADECYLTRIMETHYLAMMONIUM"/CN
	1	"2-PYRIDINOL, 1-OXIDE, LITHIUM SALT"/CN
	1	"2-PYRIDINOL, 1-OXIDE, POTASSIUM SALT"/CN
	1	"2-PYRIDINOL, 1-OXIDE, SODIUM SALT"/CN
	1	"2-PYRIDINOL, 1-OXIDE, ZINC SALT"/CN
L1	9	("2-PYRIDINOL, 1-OXIDE"/CN OR "2-PYRIDINOL, 1-OXIDE, COMPD. WITH DODECYLAMINE"/CN OR "2-PYRIDINOL, 1-OXIDE, COMPD. WITH DODECYLAMINE (1:1)"/CN OR "2-PYRIDINOL, 1-OXIDE, ION(1-)/CN OR

"2-PYRIDINOL, 1-OXIDE, ION(1-), HEXADECYLTRIMETHYLAMMONIUM"/CN  
OR "2-PYRIDINOL, 1-OXIDE, LITHIUM SALT"/CN OR "2-PYRIDINOL,  
1-OXIDE, POTASSIUM SALT"/CN OR "2-PYRIDINOL, 1-OXIDE, SODIUM  
SALT"/CN OR "2-PYRIDINOL, 1-OXIDE, ZINC SALT"/CN)

=> fil .genbiotech

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	34.82	34.97

FILE 'MEDLINE' ENTERED AT 08:01:00 ON 18 JUN 2001

FILE 'CANCERLIT' ENTERED AT 08:01:00 ON 18 JUN 2001

FILE 'BIOSIS' ENTERED AT 08:01:00 ON 18 JUN 2001  
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'CAPLUS' ENTERED AT 08:01:00 ON 18 JUN 2001  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOTECHDS' ENTERED AT 08:01:00 ON 18 JUN 2001  
COPYRIGHT (C) 2001 DERWENT INFORMATION LTD

FILE 'EMBASE' ENTERED AT 08:01:00 ON 18 JUN 2001  
COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'WPIDS' ENTERED AT 08:01:00 ON 18 JUN 2001  
COPYRIGHT (C) 2001 DERWENT INFORMATION LTD

=> s 11

EXCEEDS MAXIMUM FIELD LENGTH, WILL BE SEARCHED AS '2-PYRIDINOL, 1-OXIDE, ION(1-),  
, HEXADECYLTRIMETHYLAMMO/CN'

EXCEEDS MAXIMUM FIELD LENGTH, WILL BE SEARCHED AS '2-PYRIDINOL, 1-OXIDE, ION(1-),  
, HEXADECYLTRIMETHYLAMMO/CN'

L2 67 L1

=> s (2-pyridinol or 2-hydroxypyridine) (a) (n-oxide or 1-oxide)

4 FILES SEARCHED...

L3 121 (2-PYRIDINOL OR 2-HYDROXYPYRIDINE) (A) (N-OXIDE OR 1-OXIDE)

=> s (2-pyridinol or 2-hydroxypyridine) (w) (n-oxide or 1-oxide)

4 FILES SEARCHED...

L4 117 (2-PYRIDINOL OR 2-HYDROXYPYRIDINE) (W) (N-OXIDE OR 1-OXIDE)

=> s 12 or 13

L5 146 L2 OR L3

=> dup rem 15

PROCESSING COMPLETED FOR L5

L6 117 DUP REM L5 (29 DUPLICATES REMOVED)

=> s l6 not (py=2000 or py=2001)

L7 101 L6 NOT (PY=2000 OR PY=2001)

=> d l7 ti 1-

YOU HAVE REQUESTED DATA FROM 101 ANSWERS - CONTINUE? Y/(N):y

L7 ANSWER 1 OF 101 MEDLINE

TI Comparison of the effects of pyridine and its metabolites on rat liver and kidney.

L7 ANSWER 2 OF 101 MEDLINE

TI Chelators affecting iron absorption in mice.

L7 ANSWER 3 OF 101 MEDLINE

TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxides\*\*\* :  
effective new chelators in iron mobilisation.

L7 ANSWER 4 OF 101 MEDLINE

TI Iron mobilisation from lactoferrin by chelators at physiological pH.

L7 ANSWER 5 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

TI Substrate, substrate analogue and inhibitor interactions with the ferrous active site of catechol 2,3-dioxygenase monitored through XAS studies.

L7 ANSWER 6 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

TI Vanadium(IV) and oxovanadium(IV) complexes of hydroxamic acids and related ligands.

L7 ANSWER 7 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

TI INTERACTIONS OF \*\*\*2\*\*\* \*\*\*HYDROXYPYRIDINE\*\*\* \*\*\*N\*\*\* -  
\*\*\*OXIDE\*\*\* WITH BIOLOGICAL CATIONS CALCIUM MAGNESIUM ZINC MANGANESE.

L7 ANSWER 8 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

TI PROTO CATECHUATE 3 4 DI OXYGENASE EC-1.13.11.3 COMPARATIVE STUDY OF  
INHIBITION AND ACTIVE SITE INTERACTIONS OF PYRIDINE N OXIDES.

L7 ANSWER 9 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

TI PROTO CATECHUATE 3 4 DI OXYGENASE MECHANISTIC IMPLICATIONS OF INHIBITION  
BY THE TRANSITION STATE ANALOG 2 HYDROXY ISO NICOTINIC-ACID N OXIDE.

L7 ANSWER 10 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

TI STUDIES ON 1 3 BENZOXAZINES 3. REACTION OF IMIDOYL CHLORIDES OF 1 3  
BENZOXAZINES WITH 2 HYDROXY PYRIDINE N OXIDE OR 2 MERCAPTO PYRIDINE N  
OXIDE A NOVEL SULFUR NITROGEN BOND FORMATION VIA ELECTRO CYCLIC  
REARRANGEMENT.

L7 ANSWER 11 OF 101 CAPLUS COPYRIGHT 2001 ACS

TI Towards the reaction mechanism of pyrogallol-phloroglucinol  
transhydroxylase of Pelobacter acidigallici

L7 ANSWER 12 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Equilibrium studies on copper(II)- and iron(III)- monohydroxamates

L7 ANSWER 13 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Alkyl transfer with retention and inversion of configuration:  
reexamination of a putative [1s,4s] sigmatropic rearrangement

L7 ANSWER 14 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Heat-developable image-recording element

L7 ANSWER 15 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Image-recording element

L7 ANSWER 16 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Liquid propellant XM46 stability enhancement

L7 ANSWER 17 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Laundry pretreatment with peroxide bleaches containing chelators for iron,  
copper or manganese for reduced fabric damage

L7 ANSWER 18 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Photochemistry of N-Hydroxy-2(1H)-pyridone, a More Selective Source of  
Hydroxyl Radicals Than N-Hydroxypyridine-2(1H)-thione

L7 ANSWER 19 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Thiapyran formation via an unexpected thioaldehyde intermediate by the  
thermal decomposition of phenacyl sulfoxides bearing some heterocycles

L7 ANSWER 20 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Carbodiimide-Mediated Amide Formation in a Two-Phase System. A High-Yield  
and Low-Racemization Procedure for Peptide Synthesis

L7 ANSWER 21 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Structure and kinetics of pyridine N-oxide thermal degradation

L7 ANSWER 22 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Structure-activity relationship of ligands of uracil  
phosphoribosyltransferase from *Toxoplasma gondii*

L7 ANSWER 23 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Minoxidil analogs as inhibitors of cell proliferation and lysyl  
hydroxylase

L7 ANSWER 24 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI An ab initio study of the tautomeric equilibria of the N-oxides of  
hydroxypyridines in the vapor phase

L7 ANSWER 25 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Synergistic wood preservative compositions comprising phenol and  
pyrithione derivatives

L7 ANSWER 26 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Preservation of diagnostic test reagents and kits

L7 ANSWER 27 OF 101 CAPLUS COPYRIGHT 2001 ACS  
TI Cosmetic preservative reference materials

L7 ANSWER 28 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Structure-activity relationship of minoxidil analogs as inhibitors of lysyl hydroxylase in cultured fibroblasts

L7 ANSWER 29 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Interactions of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\* with biological cations (calcium(2+), magnesium(2+), zinc(2+) manganese(2+)...)

L7 ANSWER 30 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI A proton and carbon-13 NMR and x-ray diffraction study of the tautomerism of 2-hydroxy- and 2,3-dihydroxypyridine N-oxides. X-ray molecular structure of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\*

L7 ANSWER 31 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Wood preservatives

L7 ANSWER 32 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Synergistic wood preservatives comprising quaternary ammonium compounds and pyridine N-oxide derivatives

L7 ANSWER 33 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Iron chelators

L7 ANSWER 34 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Free radical and cytotoxic effects of chelators and their iron complexes in the hepatocyte

L7 ANSWER 35 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Mobilization of plutonium and iron from transferrin and ferritin by hydroxypyridone chelators

L7 ANSWER 36 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Molecular orbital study of some aromatic N-oxide systems

L7 ANSWER 37 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Ligands for metalizable dyes

L7 ANSWER 38 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Isomeric methoxypyridine 1-oxides and 1-methoxypyridones: electronic spectra and structure

L7 ANSWER 39 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI [170]water and nitric oxide binding by protococatechuate 4,5-dioxygenase and catechol 2,3-dioxygenase. Evidence for binding of exogenous ligands to the active site iron of extradiol dioxygenases

L7 ANSWER 40 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Pyridine oxide derivative pharmaceuticals and cosmetics with reduced toxicity

L7 ANSWER 41 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Feed compositions containing copper salts of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxides\*\*\*

L7 ANSWER 42 OF 101 CAPLUS COPYRIGHT 2001 ACS

TI Lanthanide and actinide complexes with bidentate ligands. Crystal structure of dimethylformamidetetrakis(1-oxo-2-thiopyridinato)thorium(IV)

L7 ANSWER 43 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI The reduction of aliphatic and aromatic N-oxides to the corresponding amines with titanium(III) chloride

L7 ANSWER 44 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Lanthanide and actinide complexes with bidentate ligands. Crystal structure of dimethylformamidetetrakis(1-oxo-2-thiopyridinato)thorium(IV)

L7 ANSWER 45 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Structural studies of organoboron compounds. XVIII. Crystal and molecular structures of 2,2-diphenyl-1,3-dioxo-3a-azonia-2-borataindan and 4-cyclohexyl-6-methyl-2,2-diphenyl-1,3-dioxo-3a-azonia-2-borataindan

L7 ANSWER 46 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Uranyl complexes with bidentate derivatives of pyridine N-oxide. The crystal structure of (dimethyl sulfoxide)bis(1-oxo-2-thiopyridinato)dioxouranium(VI) and aquabis(1,2-dioxopyridinato)dioxouranium(VI) monohydrate

L7 ANSWER 47 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Boron chelates of 2-hydroxy- and 2-mercaptopyridine N-oxides

L7 ANSWER 48 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Softening agent and treatment of textiles after washing

L7 ANSWER 49 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Metal complexes of the 2-hydroxy derivative of pyridine N-oxide

L7 ANSWER 50 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Photoelectron-determined core binding energies and predicted gas-phase basicities for the 2-hydroxypyridine .dblarw. 2-pyridone system

L7 ANSWER 51 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Isolation and characterization of a silicon-organic complex from plants

L7 ANSWER 52 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Dialkyltin salts of 2-(mercapto or hydroxy)pyridine oxide

L7 ANSWER 53 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Nitrogen-14 nuclear magnetic resonance of some monosubstituted pyridine N-oxides

L7 ANSWER 54 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Bonding state of penta- and hexacoordinate silicon in cationic and anionic o-arylenedioxy chelates

L7 ANSWER 55 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Dialkyltin salts of substituted pyridine-1-oxides

L7 ANSWER 56 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Dialkyltin salts of substituted pyridine 1-oxides as fungicides and bactericides

L7 ANSWER 57 OF 101 CAPLUS COPYRIGHT 2001 ACS

TI Polyhaloaromatic compounds. XVIII. Grignard reactions on pentachloropyridine 1-oxide

L7 ANSWER 58 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Cyano- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\*

L7 ANSWER 59 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Substitutes 2-hydroxy-5-nitropyridine 1-oxides

L7 ANSWER 60 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\*  
 derivatives

L7 ANSWER 61 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 2-Hydroxy-4-acylaminopyridine 1-oxides

L7 ANSWER 62 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Alkoxy- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxides\*\*\*

L7 ANSWER 63 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Carbamoyl 1- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* -  
 \*\*\*oxide\*\*\*

L7 ANSWER 64 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Amino- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\*

L7 ANSWER 65 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Alkoxy- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\*

L7 ANSWER 66 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI 4-Ethoxycarbonyl 1- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* -  
 \*\*\*oxide\*\*\*

L7 ANSWER 67 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Establishment and treatment of cutaneous Candida albicans infection in the rabbit

L7 ANSWER 68 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Fungicidal pharmaceutical compositions

L7 ANSWER 69 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Hydroxypyridine oxides as antifungal agents

L7 ANSWER 70 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Studies on hydrogenolysis. XLV. Selectivity of catalysts and effect of additives. 12. Selectivity of catalysts in hydrogenation of pyridine derivatives

L7 ANSWER 71 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Separation of the Si complex of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\*  
 \*\*\*N\*\*\* - \*\*\*oxide\*\*\* into optical antipodes. An evidence for the octahedral coordination around Si

L7 ANSWER 72 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Structure and behavior of organic analytical reagents. Formation constants of transition metal complexes of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\*  
 \*\*\*1\*\*\* - \*\*\*oxide\*\*\* and 2-mercaptopyridine 1-oxide



L7 ANSWER 73 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Water resistant cationic silicon complexes of \*\*\*2\*\*\* -  
 \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\*

L7 ANSWER 74 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Glycosides from heterocycles. II. Glucosides from hydroxypyridine-N-oxides

L7 ANSWER 75 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI A possible mode of action of some antifungal and antibacterial chelating agents

L7 ANSWER 76 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI N-Oxides and related compounds. V. The tautomerism of 2- and 4-amino- and -hydroxy pyridine 1-oxide

L7 ANSWER 77 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Bromination and nitration of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\*  
 \*\*\*N\*\*\* - \*\*\*oxide\*\*\*

L7 ANSWER 78 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\* and homologs

L7 ANSWER 79 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI The directive influence of the N-oxide group during the nitration of derivatives of pyridine N-oxide. III. The nitration of 2- and 3-methoxypyridine N-oxide

L7 ANSWER 80 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\* and homologs

L7 ANSWER 81 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Aspergillic acid and the chemistry of pyrazine derivatives

L7 ANSWER 82 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Hydroxamic acids. II. The synthesis and structure of cyclic hydroxamic acids from pyridine and quinoline

L7 ANSWER 83 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 TI Hydroxamic acids. I. Cyclic hydroxamic acids derived from pyridine and quinoline

L7 ANSWER 84 OF 101 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
 TI New orally active iron chelators.

L7 ANSWER 85 OF 101 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
 TI [Ligation of the internal maxillary artery in a case of epistaxis uncontrollable by the habitual methods].  
 LIGADURA DA ARTERIA MAXILAR INTERNA EM UM CASO DE EPISTAXE INCONTROLAVEL PELOS METODOS HABITUAIS.

L7 ANSWER 86 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Fungicidal water soluble wood preservative - contg. salt of 2-hydroxy pyridine, amino cpd. and copper salt.

L7 ANSWER 87 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Synergistic wood preservative compsns. - contg. 2-mercapto- or  
 2-hydroxy-pyridine N-oxide and quat. ammonium-salts.

L7 ANSWER 88 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI 1-Hydroxy pyrid-2-one derivs. - used for removal of excess body iron or  
 treatment of anaemia.

L7 ANSWER 89 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Sodium and zinc pyrithione prodn. by reacting sodium hydrosulphide - and  
 base with 2-halo pyridine-N-oxide then heating and opt. reacting with zinc  
 salt.

L7 ANSWER 90 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Synthesis of 4 substd 2 hydroxy 5 nitro pyridene 1 oxides - having  
 antimicrobial activity.

L7 ANSWER 91 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI 4-cyano-2-hydroxypiridine 1-oxide antifungal agent.

L7 ANSWER 92 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* - \*\*\*oxide\*\*\* derivs  
 antibacterial.

L7 ANSWER 93 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI 4-acetamido- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* -  
 \*\*\*oxide\*\*\* preparation antimicrobial.

L7 ANSWER 94 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* - \*\*\*oxide\*\*\* derivs  
 antibacterial.

L7 ANSWER 95 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI 4-acetamido- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* -  
 \*\*\*oxide\*\*\* preparation antimicrobial.

L7 ANSWER 96 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI 4-alkoxy- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxides\*\*\*  
 antibacterials.

L7 ANSWER 97 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI 4-alkoxycarbonyl- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* -  
 \*\*\*oxides\*\*\* .

L7 ANSWER 98 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Antimicrobial 4-carbamoyl- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\*  
 - \*\*\*oxide\*\*\* .

L7 ANSWER 99 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Antibacterial 4-alkoxy- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* -  
 \*\*\*oxide\*\*\* .

L7 ANSWER 100 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
 TI Antibacterial 4-amino- \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* -  
 \*\*\*oxide\*\*\* .

L7 ANSWER 101 OF 101 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD

TI Fungicidal \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* -  
\*\*\*oxides\*\*\* .

=>

=> s 17 bib abs 1-7,23,29,33,34,40,49,60,74,80

MISSING OPERATOR L7 BIB

The search profile that was entered contains terms or  
nested terms that are not separated by a logical operator.

=> d 17 bib abs 1-7,23,29,33,34,40,49,60,74,80

L7 ANSWER 1 OF 101 MEDLINE

AN 96234254 MEDLINE

DN 96234254 PubMed ID: 8644130

TI Comparison of the effects of pyridine and its metabolites on rat liver and  
kidney.

AU Carlson G P

CS School of Health Sciences, Purdue University, West Lafayette, IN  
47907-1338, USA.

NC ES04362 (NIEHS)

SO TOXICOLOGY LETTERS, (1996 Jun) 85 (3) 173-8.

Journal code: VXN; 7709027. ISSN: 0378-4274.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199607

ED Entered STN: 19960726

Last Updated on STN: 19960726

Entered Medline: 19960716

AB In order to evaluate the possibility that the metabolism of pyridine may  
be important for its toxic actions, pyridine was compared with pyridine  
\*\*\*N\*\*\* - \*\*\*oxide\*\*\* , \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* ,  
3-hydroxypyridine, 4-hydroxypyridine and pyridinium methyliodide in rats  
given equal molar doses of the chemicals i.p. Hepatotoxicity was assessed by  
measuring serum sorbitol dehydrogenase, nephrotoxicity by determining  
increases in blood urea nitrogen and serum creatinine, and influence on  
xenobiotic metabolism by measuring changes in p-nitrophenol hydroxylase  
and ethoxyresorufin and benzyloxyresorufin dealkylase activities. After a  
single dose of 2.5 mmol/kg, pyridinium methyliodide was the only compound  
that was lethal whereas 2-hydroxypyridine was the only one that caused  
significant hepatotoxicity. Pyridine, pyridine N-oxide, 3-hydroxypyridine  
and 4-hydroxypyridine were effective inducers of xenobiotic metabolism.  
Thus the metabolites of pyridine may play a role, either singly or  
collectively, in the actions of pyridine.

L7 ANSWER 2 OF 101 MEDLINE

AN 91248322 MEDLINE

DN 91248322 PubMed ID: 2095129

TI Chelators affecting iron absorption in mice.

AU Kontoghiorghe G J

CS Department of Haematology, Royal Free Hospital School of Medicine, London,  
UK.

SO ARZNEIMITTEL-FORSCHUNG, (1990 Dec) 40 (12) 1332-5.  
Journal code: 91U; 0372660. ISSN: 0004-4172.

CY GERMANY: Germany, Federal Republic of

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199107

ED Entered STN: 19910719  
Last Updated on STN: 19980206  
Entered Medline: 19910702

AB The effect of natural and synthetic chelators on iron (<sup>59</sup>Fe) absorption in mice has been studied in three different experiments using single or repeated intragastric administrations of chelator iron (<sup>59</sup>Fe) complexes of different chelator doses. The amount of <sup>59</sup>Fe in whole animals, their excretions and also distribution of <sup>59</sup>Fe in blood, liver, spleen and heart was measured at one, three and eight weeks following the <sup>59</sup>Fe-chelator administrations and compared to controls which received the same amount of iron (<sup>59</sup>Fe) but no chelator. 2-Hydroxy-4-methoxypyridine-1-oxide and maltol, which form lipophilic iron complexes, were found to cause an increase of <sup>59</sup>Fe absorption while other chelators caused a decrease either by precipitating iron eg. \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*1\*\*\* - \*\*\*oxide\*\*\* or by forming non absorbable soluble iron complexes eg. desferrioxamine, mimosine, EDTA. 1,2-Dimethyl-3-hydroxypyrid-4-one caused a decrease in iron absorption at a high dose (10 mg) by comparison to the control group but it did not significantly alter iron absorption at a lower dose (2 mg). It is suggested that natural and synthetic iron chelating compounds influence the absorption of iron and some may have a use in the treatment of diseases associated with gastro-intestinal iron absorption imbalance.

L7 ANSWER 3 OF 101 MEDLINE

AN 87157804 MEDLINE

DN 87157804 PubMed ID: 3828392

TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxides\*\*\* :  
effective new chelators in iron mobilisation.

AU Kontoghiorghes G J

SO BIOCHIMICA ET BIOPHYSICA ACTA, (1987 Apr 16) 924 (1) 13-8.  
Journal code: A0W; 0217513. ISSN: 0006-3002.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 198705

ED Entered STN: 19900303  
Last Updated on STN: 19970203  
Entered Medline: 19870521

AB The \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxide\*\*\* derivatives, \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxide\*\*\*, 2,4-dihydroxypyridine-N-oxide, 2-hydroxy-4-methoxypyridine-N-oxide and 2-hydroxy-4-(2'-methoxyethoxy)pyridine-N-oxide have been shown to remove iron from human transferrin and horse spleen ferritin at pH 7.4 at levels higher than those caused by desferrioxamine. Their reactions with transferrin were mainly biphasic and took 2-5 h to reach completion but iron mobilisation from ferritin was slower and their reactions continued after 40 h of incubation. The intraperitoneal and intragastric administration of 2,4-dihydroxypyridine-N-oxide to two iron-loaded

59Fe-labelled mice caused an increase in 59Fe excretion which is comparable to that caused by desferrioxamine intraperitoneally. These results increase the prospects for the use of these chelators as probes for studying iron metabolism and in the treatment of iron overload and other diseases of iron imbalance.

- L7 ANSWER 4 OF 101 MEDLINE  
AN 86216301 MEDLINE  
DN 86216301 PubMed ID: 3708002  
TI Iron mobilisation from lactoferrin by chelators at physiological pH.  
AU Kontoghiorghes G J  
SO BIOCHIMICA ET BIOPHYSICA ACTA, (1986 Jun 19) 882 (2) 267-70.  
Journal code: AOW; 0217513. ISSN: 0006-3002.  
CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 198607  
ED Entered STN: 19900321  
Last Updated on STN: 19970203  
Entered Medline: 19860724  
AB Several alpha-ketohydroxypyridine, \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\*  
\*\*\*N\*\*\* - \*\*\*oxide\*\*\* and 8-hydroxyquinoline chelators were shown to mobilise iron from diferric 59Fe-labelled human lactoferrin at physiological pH without the use of mediators or reducing agents. 1,2-Dimethyl-3-hydroxypyrid-4-one was found to be the most effective chelator, removing 90% of 59Fe from [59Fe]lactoferrin, in contrast to desferrioxamine, which was ineffective under the same conditions.
- L7 ANSWER 5 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS  
AN 1994:504817 BIOSIS  
DN PREV199497517817  
TI Substrate, substrate analogue and inhibitor interactions with the ferrous active site of catechol 2,3-dioxygenase monitored through XAS studies.  
AU Bertini Vvano, Fabrizio Briganti; Mangani, Stefano; Nolting, Hans F.; Scozzafava, Andrea (1)  
CS (1) Dip. Chim., Univ. Firenze, Via Gino Capponi 7, I-50121 Firenze Italy  
SO FEBS Letters, (1994) Vol. 350, No. 2-3, pp. 207-212.  
ISSN: 0014-5793.  
DT Article  
LA English  
AB The interactions of catechol (substrate). 2-hydroxy-pyridine-N-oxide (substrate analogue), and 2-bromophenol (inhibitor) with the extradiol cleaving catechol-2,3-dioxygenase from Pseudomonas putida mt-2 have been monitored through X-ray absorption spectroscopy (XAS). The analysis of the data provides details about the mode of coordination of the substrate and of the inhibitors to the active site of the enzyme.
- L7 ANSWER 6 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS  
AN 1993:160778 BIOSIS  
DN PREV199395081828  
TI Vanadium(IV) and oxovanadium(IV) complexes of hydroxamic acids and related ligands.  
AU Dessi, Alessandro; Micera, Giovanni (1); Sanna, Daniele; Erre, Liliana Strinna  
CS (1) Dep. Chem., Univ. Sassari, Via Vienna 2, 07100 Sassari Italy  
SO Journal of Inorganic Biochemistry, (1992) Vol. 48, No. 4, pp. 279-287.

ISSN: 0162-0134.

DT Article

LA English

AB Complex formation between oxovanadium(IV) and a series of hydroxamic or hydroxamic-like ligands (acetohydroxamic, benzohydroxamic, \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxide\*\*\* and N-phenyl-benzohydroxamic acids) has been investigated in solution by EPR spectroscopy. VO(IV) complexes involving the ligand-to-metal molar ratio of 2:1 have also been isolated in the solid state. The results show that the simple hydroxamic ligands (acetohydroxamic and benzohydroxamic acids) can undergo two deprotonation processes and thus act as either hydroxamato(1-) or hydroximato(2-) ligands. A similar dissociation pattern is not possible for N-phenyl-benzohydroxamic acid and the hydroxamic-like ligand \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* - \*\*\*N\*\*\* - \*\*\*oxide\*\*\*, which are able to yield only a hydroxamato(1-) coordinating anion. All the ligands form hexa-coordinate tris-chelated complexes of V(IV) after displacing the oxo group from VO(IV).

L7 ANSWER 7 OF 101 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1990:484493 BIOSIS

DN BR39:108514

TI INTERACTIONS OF \*\*\*2\*\*\* \*\*\*HYDROXYPYRIDINE\*\*\* \*\*\*N\*\*\* - \*\*\*OXIDE\*\*\* WITH BIOLOGICAL CATIONS CALCIUM MAGNESIUM ZINC MANGANESE.

AU DEIDA M F; PIERRARD J C; RIMBAULT J

CS UNIV. DE REIMS CHAMPAGNE-ARDENNE, FAC. DES SCI., LAB. DE CHIMIE MINERALE, BP 347, 51062 REIMS CEDEX, FR.

SO FIRST INTERNATIONAL SYMPOSIUM ON METAL IONS IN BIOLOGY AND MEDICINE, MAY 16-19, 1990. TRACE ELEM MED. (1990) 7 (2), 101.

CODEN: TEMDE6. ISSN: 0174-7371.

DT Conference

FS BR; OLD

LA English

L7 ANSWER 23 OF 101 CAPLUS COPYRIGHT 2001 ACS

AN 1994:570582 CAPLUS

DN 121:170582

TI Minoxidil analogs as inhibitors of cell proliferation and lysyl hydroxylase

IN Murad, Saood; Pinnell, Sheldon R.

PA Duke University, USA

SO U.S., 15 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 5328913	A	19940712	US 1992-987952	19921211
AB	A method of inhibiting cell proliferation and lysyl hydroxylase expression by using minoxidil derivs. and analogs is decribed. The inhibitory activity of hydroxy derivs. of minoxidil is such that these compds. can be used as selective antifibrotic agents. Minoxidil and some analogs inhibited the activity of lysyl hydroxylase comparable to the parent drug and some others did not.				

L7 ANSWER 29 OF 101 CAPLUS COPYRIGHT 2001 ACS

AN 1991:215665 CAPLUS

DN 114:215665  
 TI Interactions of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* -  
 \*\*\*oxide\*\*\* with biological cations (calcium(2+), magnesium(2+),  
 zinc(2+) manganese(2+)...)

AU Deida, M. F.; Pierrard, J. C.; Rimbault, J.  
 CS Fac. Sci., Univ. Reims, Reims, 51062, Fr.  
 SO Met. Ions Biol. Med., Proc. Int. Symp., 1st (1990), 567-9. Editor(s):  
 Collery, Philippe. Publisher: Libbey, Paris, Fr.  
 CODEN: 56ZJAL

DT Conference  
 LA English

AB Coordination of Ca, Mg, Mn, Zn and Ni with the title ligand was studied at  
 25.degree. in 1M (NaClO4) soln. by a potentiometric method. Stability  
 consts. of the complexes are given. The order of stability of the  
 complexes vary with the nature of the metal ion as follows:  $\text{Ca}^{2+} < \text{Mg}^{2+} < \text{Mn}^{2+} < \text{Zn}^{2+} \approx \text{Ni}^{2+}$ . The ionization const. of the ligand was also  
 calcd.

L7 ANSWER 33 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 AN 1989:108140 CAPLUS  
 DN 110:108140  
 TI Iron chelators

AU Mostert, L. J.; Koster, J. F.; Van Eijk, H. G.  
 CS Med. Fac., Erasmus Univ., Rotterdam, 300 DR, Neth.  
 SO Tijdschr. Ned. Ver. Klin. Chem. (1988), 13(6), 211-14  
 CODEN: TNVCE9; ISSN: 0168-8472

DT Journal  
 LA Dutch

AB In a comparative screening study of chelators intended for clin. use, 11  
 Fe chelators have been tested for their ability to mobilize  $^{59}\text{Fe}$  from  
 $^{59}\text{Fe}$ -labeled ferritin and from rat hepatocytes labeled with  
 $^{59}\text{Fe}$ -transferrin. The toxic effects of the chelators were also studied  
 using microsomal lipid peroxidn. induced by  $\text{Fe}^{3+}/\text{ADP}$  and NADPH. Mimosine  
 and 1,2-dimethyl-3-hydroxypyridin-4-one were the most effective chelators  
 in iron mobilization and did not catalyze lipid peroxidn. Besides  
 investigating the iron-binding capacity of new chelators, their ability to  
 catalyze lipid peroxidn. has to be ruled out.

L7 ANSWER 34 OF 101 CAPLUS COPYRIGHT 2001 ACS  
 AN 1987:629106 CAPLUS  
 DN 107:229106  
 TI Free radical and cytotoxic effects of chelators and their iron complexes  
 in the hepatocyte

AU Mostert, L. J.; Van Dorst, J. A. L. M.; Koster, J. F.; Van Eijk, H. G.;  
 Kontoghiorghe, G. J.  
 CS Dep. Chem. Pathol. Biochem. I, Erasmus Univ., Rotterdam, Neth.  
 SO Free Radical Res. Commun. (1987), 3(6), 379-88  
 CODEN: FRRCEX; ISSN: 8755-0199

DT Journal  
 LA English

AB In a comparative screening study of chelators intended for clin. use  
 eleven iron chelators have been tested for their ability to mobilize  $^{59}\text{Fe}$   
 from  $^{59}\text{Fe}$ -labeled ferritin and from hepatocytes of rats labeled with  
 $^{59}\text{Fe}$ -transferrin. The toxic effects of the chelators were also studied  
 using microsomal lipid peroxidn. induced by  $\text{Fe}^{3+}/\text{ADP}$  and NADPH. From  
 these tests it was shown that 1,2-di-Me 3-hydroxypyrid-4-one (L1) and  
 mimosine were the most effective Fe chelators in Fe mobilization and did

not catalyze lipid peroxidn. Thus, aside from their Fe binding properties, chelators should be examd. for their role in catalyzing lipid peroxidn. in toxicol. screening.

L7 ANSWER 40 OF 101 CAPLUS COPYRIGHT 2001 ACS  
AN 1985:172662 CAPLUS  
DN 102:172662  
TI Pyridine oxide deriative pharmaceuticals and cosmetics with reduced toxicity  
IN Gari, Kailash Kumar  
PA Paninkret Chemisch Pharmazeutisches Werk G.m.b.H., Fed. Rep. Ger.  
SO Ger. Offen., 18 pp.  
CODEN: GWXXBX  
DT Patent  
LA German  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 3327485	A1	19850214	DE 1983-3327485	19830729
AB	The toxicity of cosmetics and pharmaceuticals contg. 2-mercaptopyridine N-oxide [1121-31-9] or ***2*** - ***hydroxypyridine*** ***N*** - ***oxide*** [ ***13161-30-3*** ] or their salts is decreased by the addn. of a proteolyzate of plant or animal origin at a ratio of 0.5-20 parts proteolyzate to 1 part pyridine oxide. Thus, slaughterhouse coagulated blood was sepd. from plasma, dild. with H2O to 10% solids, and mixed with equal parts of pancreatin, papain, and 2 yeast enzymes (10 g/10 L) for 1 h at 35-38.degree., held at 53-57.degree. for 2 h, mixed with a filter aid, heated to 90.degree., and filtered. The filtrate was concd. and spray-dried, suspended in 70% EtOH, and filtered after 8-12 h. EtOH was evapd., and the residue was spray-dried, and optionally dialyzed. A soln. for topical use was prepd. from 100 g Na 2-mercaptopyridine N-oxide [15922-78-8] and 600 g of the blood hydrolyzate in 1 L 60% iso-PrOH. The effectiveness of the hydrolyzate in decreasing the retinal toxicity of Na pyridinethione was demonstrated by electroretinogram b wave potentials in dark-adapted rats injected i.p. with the compd. and the hydrolyzate.				

L7 ANSWER 49 OF 101 CAPLUS COPYRIGHT 2001 ACS  
AN 1981:596537 CAPLUS  
DN 95:196537  
TI Metal complexes of the 2-hydroxy derivative of pyridine N-oxide  
AU Landers, Arthur E.; Phillips, David J.  
CS Sch. Chem., Univ. New South Wales, Kensington, 2033, Australia  
SO Inorg. Chim. Acta (1981), 51(1), 109-15  
CODEN: ICHAA3; ISSN: 0020-1693

DT Journal  
LA English  
AB Cu(opo)2 (Hopo = 1-hydroxy-2-pyridone, the tautomeric form of \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\* ), Cu(opo)NO3.H2O, Cu(opo)Br.0.5MeOH, Cu(opo)Cl, Ni(opo)Cl.1.5H2O, Co(opo)Cl.1.5H2O, and Fe(opo)3.H2O were prepd. Structures with bidentate 1-hydroxy-2-pyridone are proposed for the complexes on the basis of measurements of x-ray powder diffraction spectra. Moessbauer, IR and electronic spectra, and magnetic data at >89 K are reported. Cu(opo)Br.0.5MeOH is antiferromagnetic, and its susceptibility data was fitted to various models.

L7 ANSWER 60 OF 101 CAPLUS COPYRIGHT 2001 ACS



AN 1970:12596 CAPLUS  
 DN 72:12596  
 TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\*  
 derivatives  
 IN Minakami, Satoshi; Hirai, Eizo  
 PA Shionogi and Co., Ltd.  
 SO Japan., 2 pp.  
 CODEN: JAXXAD  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 44025582	B4	19691028	JP	19650302
GI	For diagram(s), see printed CA Issue.				
AB	Manuf. of I, useful as a bactericide, is described. In an example, 2.5 g 4-chloro-2-methoxypyridine 1-oxide is treated with 75 ml AcCl, the mixt. evapd. in vacuo, the residual oil heated with 100 ml H2O and 4 ml concd. HCl 10 hr, concd. in vacuo, and cooled with ice to give 9 g I (R = Cl), m. 145-6.degree. (H2O). Similarly prepd. are the following I (R and m.p. given): NO2, 169-71.degree.; CO2H, >250.degree..				

L7 ANSWER 74 OF 101 CAPLUS COPYRIGHT 2001 ACS

AN 1963:409283 CAPLUS

DN 59:9283

OREF 59:1744c-g

TI Glycosides from heterocycles. II. Glucosides from hydroxypyridine-N-oxides

AU Wagner, G.; Pischel, H.

CS Karl-Marx Univ., Leipzig, Germany

SO Arch. Pharm. (1962), 295, 897-910

DT Journal

LA Unavailable

AB cf. CA 58, 4642b. The glucosides of the tautomers of 2- and 4-hydroxypyridines were prepd., as well as that from the 3-isomer. 3-Hydroxypyridine tetra-O-acetyl-.beta.-D-glucopyranoside was converted to the N-oxide with PhCO2OH, m. 78-80.degree. (HOAc-petr. ether) [.alpha.]24D -40.7 (c 5.1, CHCl3), and deacetylated with NaOMe in MeOH, m. 212.5-13.5.degree. (decompn.), [.alpha.]22D -93.8.degree. (c 2.6, H2O). Similarly were prepd. 4-hydroxypyridine N-oxide tetra-O-acetyl-.beta.-D-glucopyranoside, [.alpha.]22D -26.7.degree. (c 2.5, H2O); and 4-hydroxypyridine N-oxide .beta.-D-glucopyranoside, m. 195-7.degree., [.alpha.]21D -99.4.degree. (c 1.0, H2O). Ag salt of N-hydroxy-2-pyridone (1.6 g.) and 3.0 g. (I) tetra-O-acetyl-.alpha.-D-glucopyranosyl bromide (I) was boiled in 40 ml. dry PhMe 8 min., filtered while hot, and C6H6 added to give a white powder, recrystd. from HOAc-petr. ether, m. 142-3.degree., [.alpha.]22D -132.4.degree. (c 5.0, CHCl3), N-hydroxy-2-pyridone tetra-O-acetyl-.beta.-D-glucopyranoside (II). The Ag salt was obtained from \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\* in aq. NaOH with AgNO3. 2-Bromopyridine N-oxide (7 g.) was heated on a water bath with 50 ml. 10% NaOH 90 min., cooled, acidified with 30% H2SO4 (Congo), evapd. to dryness, the residue extd. with CHCl3, and the solvent removed to give 80% crude \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\*, m. 148-9.degree. (EtOAc). II was also prepd. from \*\*\*2\*\*\* - \*\*\*hydroxypyridine\*\*\* \*\*\*N\*\*\* - \*\*\*oxide\*\*\*, KOH, and I; and I and 2-ethoxypyridine N-oxide.

II sapond. with NaOMe gave 65% N-hydroxy-2-pyridone .beta.-D-

glucopyranoside, m. 195.5-6.5.degree. (PrOH), [.alpha.]20D -100.3.degree. (c 2.5, H2O). The Ag salt of 3-hydroxypyridine N-oxide and I in MePh gave a mixt. of 3-hydroxypyridine N-oxide tetra-O-acetyl-.beta.-D-glucopyranoside and the corresponding de-N-oxide. Treatment of the Ag salt from 4-hydroxypyridine N-oxide with I in PhMe gave a mixt. of the two tautomeric derivs. 4-N-hydroxypyridone 4-tetra-O-acetyl-.beta.-D-glucopyranoside (III) could be isolated from the mixt. by addn. of petr.-ether, soln. in CHCl3, washing with Na2CO3, dropping into petr.-ether, and recrystn. from EtOAc-petr. ether, m. 134-6.degree., [.alpha.]19D -26.2.degree. (c 2.0, CHCl3). Deacetylation gave the glucoside, m. 125-7.degree., [.alpha.]20D -33.1.degree. (c 1.0, H2O). 4-Ethoxypyridine N-oxide and I 4 days at 65.degree. gave a mixt. of III and 4-pyridone N-tetra-O-acetyl-.beta.-D-glucopyranoside (IV), identified through paper chromatography of their deacetylated glucosides. I (3 g.) and 3 g. 4-EtO-C5H5N heated in a sealed tube 4 days at 65.degree., dissolved in CHCl3, washed with HOAc, H2O, dried, evapd., and recrystd. from 50 ml. MeOH gave IV, glass, [.alpha.]20D 31.0, (c 5.0, CHCl3). Deacetylation gave an amorphous material contg. 1.5 mols. H2O, [.alpha.]19D 57.7.degree., (c 2.5, H2O). Redn. of 3- and 4-hydroxypyridine N-oxide tetra-O-acetyl-.beta.-D-glucopyranosides in MeOH-Raney-Ni gave the corresponding hydroxypyridines, m. 139-41.degree.; 110-13.degree., resp. A table of Rf values for the compds. discussed is appended.

L7 ANSWER 80 OF 101 CAPLUS COPYRIGHT 2001 ACS

AN 1951:36244 CAPLUS

DN 45:36244

OREF 45:6224a-f

TI \*\*\*2\*\*\* - \*\*\*Hydroxypyridine\*\*\* \*\*\*1\*\*\* - \*\*\*oxide\*\*\* and  
homologs

IN Shaw, Elliott N.

PA E. R. Squibb & Sons

DT Patent

LA Unavailable

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 2540218		19510206	US	
AB	<p>The compds. are antibacterial agents comparable to aspergillic acid. They occur also in the tautomeric form as 1-hydroxy-2(1H)-pyridone (I) and derivs. Reflux 2-bromopyridine 70 with PhCH2ONa 60 g. in PhCH2OH 200 cc. 2 hrs., pour into H2O, ext. with Et2O, fractionate in vacuo, and collect 38.5 g. 2-benzyloxypyridine (II), b2 134-5.degree.. Oxidize II 37 g. with 1.5 equivs. BzO2H in 700 cc. CHCl3 at room temp., wash after 3 days with NaHCO3 soln. and H2O, dry with MgSO4, evap. in vacuo, slurry the residue with EtOAc + C6H8, and filter the residue (18 g.) of 2-benzyloxypyridine 1-oxide (III), m. 102-6.degree. (from EtOH), 4.3 g. of which, boiled 10 min. with 15 cc. 20% HCl, evapd. in vacuo, and crystd. from C6H6 + MeOH gives 1.6 g. I, m. 148-50.degree. (from MeOH). Extg. the mother liquid with EtOAc gives a small quantity of an isomer of III, m. 83-4.degree.. I is also prepd. in 69% yield by shaking III with 5% Pd on charcoal in EtOH under 50 lb. H pressure. It gives a red color with FeCl3 in EtOH and forms a Cu salt, (C5H4O2N)2Cu, m. 298.degree. (decompn.). I 3 g. in AcOH 15 cc. with Br 4.3 g. in AcOH 20 ml. at room temp. during 2 hrs. gives on evapn. in vacuo 0.65 g. of a Br deriv. of I, m. 208-9.degree.. I 3.1 g. gives with HNO3 2 in AcOH 15 cc. 67% orange crystals of a nitro deriv., m. 198-9.degree.. 3-Hydroxypyridine gives with 1.5 equivs. BzO2H in CHCl3 in</p>				

12-16 hrs. at room temp. 65% 3-hydroxypyridine 1-oxide, m. 189-91.degree. (from MeOH). Refluxing 4-(1-pyridyl)pyridinium dichloride 65 with Na 13 g. in PhCH2OH 400 cc. 4 hrs., pouring into H2O, extn. with Et2O, and fractionation gives an oil, b4 147-60.degree.; redistn. gives 11.5 g. 4-benzyloxypyridine (IV), b4 155-60.degree., m. 55-6.degree.; picrate, m. 150-1.degree.. Oxidation of IV with BzO2H gives 4-benzyloxypyridine 1-oxide, m. 178-9.degree., reduced with Pd and H to 1-hydroxy-4(1H)-pyridone, m. 238-40.degree.. Carbostyryl gives with BzO2H in CHCl3 gives in 7 days on evapn., extn. with NaHCO3 soln., pptn. with HCl, and crystn. from C6H6 1-hydroxy-2(1H)-quinolone, m. 190-2.degree.. By analogous procedures are prepd. 1-hydroxy-4-methyl-2(1H)-pyridone m. 129-30.degree.; 1-hydroxy-5-bromo-2(1H)-pyridone, m. 155-7.degree.; 2-benzyloxy-5-bromopyridine 1-oxide, m. 127-8.degree.; 1-hydroxy-7-chloro-4(1H)-quinolone, m. 262.degree.; 1-hydroxy-3,6-di-sec-butyl-2(1H)-pyrazinone, from 2-chloro-3,6-di-sec-butylpyrazine.

=> fil stnguide

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	110.74	145.71
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-5.29	-5.29

FILE 'STNGUIDE' ENTERED AT 08:13:26 ON 18 JUN 2001  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.  
 LAST RELOADED: Jun 15, 2001 (20010615/UP).

=> log y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.00	145.71
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-5.29

STN INTERNATIONAL LOGOFF AT 08:26:52 ON 18 JUN 2001